arXiv:1208.4430v1 [math.AT] 22 Aug 2012

THE TOPOLOGICAL PERIOD-INDEX PROBLEM OVER 6-COMPLEXES

BENJAMIN ANTIEAU* AND BEN WILLIAMS

ABSTRACT. We completely solve the topological period-index problem for finite CW complexes of cohomological dimension at most 6. While for odd primes, the solution agrees with what is expected in algebraic geometry, the solution at the prime 2 does not, which suggests that something interesting is happening: either some restriction on the homotopy type of complex 3-folds is intervening, or the period-index conjecture is false at the prime 2 in dimension 3. At present, we cannot conclude which of these possibilities occurs.

1. INTRODUCTION

The cohomological Brauer group of a topological space X is $\mathrm{Br}'_{\mathrm{top}}(X) = \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$. Given a PU_n-torsor $P \to X$, there is a class $\delta_n(P) \in \mathrm{Br}'_{\mathrm{top}}(X)$ that is the obstruction to lifting P to a U_n-torsor; these classes arise from the exact sequences $1 \to S^1 \to \mathrm{U}_n \to \mathrm{PU}_n \to 1$. Given $\alpha \in \mathrm{Br}'_{\mathrm{top}}(X)$, the period of α, written $\mathrm{per}_{\mathrm{top}}(\alpha)$, is the order of α in the cohomological Brauer group. The index $\mathrm{ind}_{\mathrm{top}}(\alpha)$ is a more interesting invariant. It is the GCD of all n such that $\alpha = \delta_n(P)$ for some P. In general, $\mathrm{per}_{\mathrm{top}}(\alpha) | \mathrm{ind}_{\mathrm{top}}(\alpha)$. The Brauer group $\mathrm{Br}_{\mathrm{top}}(X)$ is the subgroup of $\mathrm{Br}'_{\mathrm{top}}(X)$ consisting of those classes represented by some projective bundles, i.e., those classes α for which $\mathrm{ind}_{\mathrm{top}}(\alpha)$ is finite. When X is a finite CW complex, Serre showed [11] that $\mathrm{Br}_{\mathrm{top}}(X) = \mathrm{Br}'_{\mathrm{top}}(X)$. In this case, the period and the index have the same prime divisors [1, Theorem 3.1].

There are similar definitions for schemes X. The cohomological Brauer group is $\mathrm{Br}'(X)$ is $\mathrm{H}^2_{\text{ét}}(X, \mathbb{G}_m)_{\mathrm{tors}}$. The classes $\alpha \in \mathrm{Br}'(X)$ appear as obstructions to lifting PGL_n-torsors to GL_n-torsors, and we can define $\mathrm{per}(\alpha)$ and $\mathrm{ind}(\alpha)$ in the same way. The period-index problem is to relate the period and index, and especially to find upper bounds on the index in terms of the period. The following is the major conjecture on the subject. For details, see [1].

Conjecture 1.1 (Period-index conjecture). *Let X be a d-dimensional irreducible variety over an algebraically closed field k. Then, for any $\alpha \in \mathrm{Br}(k(X))$,*

$$\mathrm{ind}(\alpha) | \mathrm{per}(\alpha)^{d-1}.$$

The conjecture has important consequences for rational points of homogeneous spaces, moduli spaces of vector bundles, quadratic forms, and division algebras. It is known from various examples, such as can be found in [8], that the exponent $d-1$ is the smallest possible. The only other significant evidence for the conjecture is the proof of the conjecture when $d = 2$. The original proof was given by de Jong [9], assuming that $\mathrm{per}(\alpha)$ was prime to the characteristic of k. Lieblich [13] gave a proof which worked in general. The conjecture is not known for a single higher-dimensional variety; even $X = \mathbb{P}^3$ remains out of reach.

In [1], we formulated the period-index problem for topological spaces, and we proved, for example, the following theorem using stable homotopy theory and twisted K-theory.

2010 *Mathematics Subject Classification.* Primary 14F22, 19L50. Secondary 55R40.

Key words and phrases. Brauer groups, twisted K-theory, period-index problems.

*The author was supported in part by the NSF under Grant RTG DMS 0838697.

Theorem 1.2. *Let X be a finite CW complex of cohomological dimension $2d$, and let $\alpha \in \mathrm{Br}'_{\mathrm{top}}(X)$ be a class such that the prime divisors of $\mathrm{per}_{\mathrm{top}}(X)$ are all greater than $2d+1$. Then,*

$$\mathrm{ind}_{\mathrm{top}}(\alpha) | \mathrm{per}_{\mathrm{top}}(\alpha)^d.$$

Our method in [1] also shows that for any fixed n, there exists an integer $e(d,n)$ such that

$$\mathrm{ind}_{\mathrm{top}}(\alpha) | \mathrm{per}_{\mathrm{top}}(\alpha)^{e(d,n)}$$

for all classes $\alpha \in \mathrm{Br}'(X)$ such that X is a finite CW complex of cohomological dimension d and $\mathrm{per}_{\mathrm{top}}(\alpha) = n$. The analogous result for function fields is unknown, except when $n = 2$ by unpublished results of Daniel Krashen.

There is as much evidence for the algebraic period-index conjecture as there is in the literature for the following guess, namely only evidence up to $d = 2$.

Statement 1.3. *Let X be a $2d$-dimensional finite CW complex, and let $\alpha \in \mathrm{Br}'_{\mathrm{top}}(X)$. Then,*

$$\mathrm{ind}_{\mathrm{top}}(\alpha) | \mathrm{per}_{\mathrm{top}}(\alpha)^{d-1}.$$

This statement is false, as we show in this paper. It fails in general for 6-dimensional complexes when 2 divides the period. It is true for 6-dimensional complexes when the period is odd. We show in the final section of the paper how this might be used to construct a counterexample to the original period-index conjecture. In light of this result, one is lead to the problem of determining the correct exponent in the topological period-index conjecture. We intend to return to this question in a sequel.

Our method in this work involves playing twisted K-theory off of some new calculations of the cohomology of $\mathrm{B\,PU}_n$. We hope that these topological results will help shed light on the algebraic period-index problem in the future.

2. RESULTS

For a natural number n we write

$$\epsilon(n) = \begin{cases} 2n & \text{if } n \text{ is even,} \\ n & \text{if } n \text{ is odd.} \end{cases} \tag{1}$$

We write β_n for the unreduced Bockstein: $\beta : \mathrm{H}^i(X, \mathbb{Z}/n) \to \mathrm{H}^{i+1}(X, \mathbb{Z})$, and we abbreviate this to β if n is clear from the context.

We will construct in Section 3 a secondary cohomology operation G defined for classes $\alpha \in \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$ and taking values in $\mathrm{H}^5(X, \mathbb{Z})/(\mathrm{H}^2(X, \mathbb{Z}) \smile \alpha)$. Any n-torsion class $\alpha \in \mathrm{H}^3(X, \mathbb{Z})$ may be written as $\beta_n(\xi)$ for some $\xi \in \mathrm{H}^2(X, \mathbb{Z}/n)$. The subgroup $\langle G(\alpha) \rangle \subset \mathrm{H}^5(X, \mathbb{Z})/(\mathrm{H}^2(X, \mathbb{Z}) \smile \alpha)$ satisfies

$$\left\langle \frac{n}{\mathrm{per}_{\mathrm{top}}(\beta_n(\xi))} G(\beta_n(\xi)) \right\rangle = \begin{cases} \langle [\beta_n(\xi^2)] \rangle & \text{if } n \text{ is odd} \\ \langle [\beta_{2n}(P_2(\xi))] \rangle & \text{if } n \text{ is even} \end{cases} \tag{2}$$

where $P_2 : \mathrm{H}^2(X, \mathbb{Z}/n) \to \mathrm{H}^4(X, \mathbb{Z}/(2n))$ is the Pontryagin square, and where the classes $[x]$ denote the reduction of $x \in \mathrm{H}^5(X, \mathbb{Z})$ to $\mathrm{H}^5(X, \mathbb{Z})/(\mathrm{H}^2(X, \mathbb{Z}) \smile \beta_n(\xi))$. The subgroup $\langle G(\alpha) \rangle$ is characterized by (2), which implies the weaker relation

$$\langle 2G(\beta_n(\xi)) \rangle = \langle [\beta_n(\xi^2)] \rangle$$

if n is even. The most useful case of equation (2) is that where $n = \mathrm{per}_{\mathrm{top}}(\beta_n(\xi))$, from which we deduce that

$$\epsilon(n) G(\alpha) = 0.$$

For any divisor m of $\epsilon(n)$, examples exist where $G(\alpha)$ has order precisely m.

Theorem A. *Suppose X is a finite CW complex of cohomological dimension $d \leq 6$, and that $\alpha \in \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$. Then*

$$\mathrm{ind}_{\mathrm{top}}(\alpha) = \mathrm{per}_{\mathrm{top}}(\alpha) \, \mathrm{ord}(G(\alpha)).$$

Theorem B. *Let X be a finite CW complex of cohomological dimension at most 6. Then, for any $\alpha \in \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$,*

$$\mathrm{ind}_{\mathrm{top}}(\alpha) \, \Bigg| \, \begin{cases} \mathrm{per}_{\mathrm{top}}(\alpha)^2 & \textit{if } \mathrm{per}_{\mathrm{top}}(\alpha) \textit{ is odd} \\ 2\,\mathrm{per}_{\mathrm{top}}(\alpha)^2 & \textit{if } \mathrm{per}_{\mathrm{top}}(\alpha) \textit{ is even.} \end{cases}$$

These upper-bounds on the index are the best possible.

As a consequence of the sharpness statement of the theorem, there exist 6-dimensional finite CW complexes X and 2-torsion classes $\alpha \in \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$ such that $2G(\alpha) \neq 0$. In light of this fact, the following theorem is especially interesting.

Theorem C. *If there exists a smooth projective complex 3-fold X and a 2-torsion class $\alpha \in \mathrm{Br}(X)$ such that $2G(\overline{\alpha}) \neq 0$, where $\overline{\alpha}$ is the image of α in $\mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$, then the period-index conjecture is false at the prime 2 in dimension 3: the image of the class α in the function field $\mathbb{C}(X)$ has period 2 and index at least 8.*

By the description of G above, to construct such a class α, it suffices to construct a smooth projective complex 3-fold X and a class $\xi \in \mathrm{H}^2(X, \mathbb{Z}/n)$ such that $\beta_2(\xi^2) \neq 0$ in $\mathrm{H}^5(X, \mathbb{Z})$.

Note that it follows by applying [10, Theorem 2.2.3] to twisted Grassmannians that if the period-index conjecture fails, then it fails for some class $\alpha \in \mathrm{Br}(X)$ where X is smooth and projective.

3. THE EXISTENCE OF G

First we show that there exists a secondary cohomology operation $G(\alpha)$ such that Theorem A holds. To do this, we recall the following three propositions. The coefficient sheaf $\mathbb{Z}(q/2)$ is $\mathbb{Z}$ if q is even and 0 if q is odd.

Proposition 3.1. *Let X be a topological space, and let $\alpha \in \mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$. Then, there is a descent spectral sequence (Atiyah-Hirzebruch spectral sequence)*

$$\mathrm{E}_2^{p,q} = \mathrm{H}^p(X, \mathbb{Z}(q/2)) \Rightarrow \mathrm{KU}^{p+q}(X)_\alpha$$

with differentials d_r^α of degree $(r, -r+1)$, which converges strongly if X is a finite dimensional CW complex. The edge map $\mathrm{KU}^0(X)_\alpha \to \mathrm{H}^0(X, \mathbb{Z})$ is the index map (or rank map, or reduced norm map).

Proof. This is [2, Theorem 4.1]. ☐

Proposition 3.2. *Let X be a connected CW complex. The differential*

$$d_3^\alpha : \mathrm{H}^0(X, \mathbb{Z}) \to \mathrm{H}^3(X, \mathbb{Z})$$

sends 1 to α.

Proof. See [3, Proposition 4.6] or [1, Proposition 2.4]. ☐

The twisted spectral sequence is a module in the natural sense over the untwisted Atiyah-Hirzebruch spectral sequence

$$\mathrm{E}_2^{p,q} = \mathrm{H}^p(X, \mathbb{Z}(q/2)) \Rightarrow \mathrm{KU}^{p+q}(X).$$

By writing a class $x \in \mathrm{H}^p(X, \mathbb{Z}(q/2))$ in the twisted spectral sequence as $x \cdot 1$, where we view 1 being twisted and x as being untwisted, we see that $d_3^\alpha(x) = x \smile d_3^\alpha(1) + d_3(x) \smile 1 = x \smile \alpha + d_3(x)$. The untwisted differential d_3 vanishes on $\mathrm{H}^2(X, \mathbb{Z})$, since there are no non-zero cohomology operations $K(\mathbb{Z}, 2) \to K(\mathbb{Z}, 5)$. Therefore, $\mathrm{E}_5^{5,-4} = \mathrm{H}^5(X, \mathbb{Z}) / (\mathrm{H}^2(X, \mathbb{Z}) \smile \alpha)$.

Proposition 3.3. *Let X be a connected finite CW complex and let $\alpha \in \mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$. Then $\mathrm{ind}_{\mathrm{top}}(\alpha)$ is the positive generator of the subgroup $\mathrm{E}_\infty^{0,0}$ of permanent cycles in $\mathrm{E}_2^{0,0} = \mathrm{H}^0(X,\mathbb{Z}) = \mathbb{Z}$ in the Atiyah-Hirzebruch spectral sequence of Proposition 3.1.*

Proof. This is an amalgamation of Proposition 2.21 and Lemma 2.23 of [1]. □

The spectral sequence of Proposition 3.1 is functorial in that a map $f : Y \to X$ and a class $\alpha \in \mathrm{H}^3(X,\mathbb{Z})$ give rise to a map of spectral sequences compatible with the map $\mathrm{KU}^*(X)_\alpha \to \mathrm{KU}^*(Y)_{f^*(\alpha)}$.

If X is a topological space, $\alpha \in \mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$ and if $\mathrm{per}_{\mathrm{top}}(\alpha) = n$, then the d_5^α-differential emanating from $\mathrm{E}_5^{0,0}$ takes the form $d_5^\alpha : n\mathbb{Z} \to \mathrm{H}^5(X,\mathbb{Z})/(\mathrm{H}^2(X,\mathbb{Z}) \smile \alpha)$. Because of the naturality of the descent spectral sequence, the assignment

$$\alpha \mapsto d_5^\alpha(n) \in \mathrm{H}^5(X,\mathbb{Z})/(\mathrm{H}^2(X,\mathbb{Z}) \smile \alpha)$$

is a secondary cohomology operation, which we denote by $G(\alpha)$.

Definition 3.4. Say that a CW complex has cohomological dimension $\leq d$ if $\mathrm{H}^n(X, A) = 0$ for all abelian sheaves A and all $n > d$.

Theorem A. *Suppose that X is a connected finite CW complex of cohomological dimension $d \leq 6$, and that $\alpha \in \mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$. Then*

$$\mathrm{ind}_{\mathrm{top}}(\alpha) = \mathrm{per}_{\mathrm{top}}(\alpha)\,\mathrm{ord}(G(\alpha)).$$

Proof. For such a space, only two nonzero differentials may emanate from $\mathrm{E}_*^{0,0}$ in the descent spectral sequence for α-twisted K-theory. The first has kernel $\mathrm{per}_{\mathrm{top}}(\alpha)\mathbb{Z}$, by Proposition 3.2 and the second has image $G(\alpha)$, and consequently kernel consisting of $\mathrm{per}_{\mathrm{top}}(\alpha)\,\mathrm{ord}(G(\alpha))\mathbb{Z} \subset \mathrm{E}_2^{0,0}$. These are the permanent cycles, and the result follows from Proposition 3.3. □

4. THE LOW-DEGREE COHOMOLOGY OF $\mathrm{B\,PU}_n$

Proposition 4.1. *For any integer $n \geq 2$, $\mathrm{H}^5(\mathrm{B\,PU}_n,\mathbb{Z}) = 0$.*

This statement can be extracted from either [12] or [16] for $\mathrm{B\,PU}_p$ when p is an odd prime. The special case of $\mathrm{B\,PU}_2$ follows from [6] and the exceptional isomorphism $\mathrm{PU}_2 \cong \mathrm{SO}_2$. We give a proof which works for all n.

To prove the proposition, we need several lemmas. Recall [15] that the ring $\mathrm{H}^*(\mathrm{BU}_n,\mathbb{Z})$ is a polynomial algebra in Chern classes $c_1, c_2, \ldots, c_n$, with $|c_i| = 2i$. Write $\rho : \mathrm{U}_n \to \mathrm{PU}_n$ for the quotient map and $\mathrm{B}\rho : \mathrm{BU}_n \to \mathrm{B\,PU}_n$ for the associated map on classifying spaces.

Lemma 4.2 (Woodward [17]). *If n is odd, then $\mathrm{B}\rho^*$ identifies $\mathrm{H}^4(\mathrm{B\,PU}_n,\mathbb{Z})$ with the subgroup of $\mathrm{H}^4(\mathrm{BU}_n,\mathbb{Z})$ generated by $\frac{n-1}{2}c_1^2 - nc_2$. If n is even, then $\mathrm{B}\rho^*$ identifies $\mathrm{H}^4(\mathrm{B\,PU}_n,\mathbb{Z})$ with the subgroup of $\mathrm{H}^4(\mathrm{BU}_n,\mathbb{Z})$ generated by $(n-1)c_1^2 - 2nc_2$.*

The proof is included for the sake of completeness.

Proof. There is a fiber sequence $\mathrm{B}S^1 \to \mathrm{BU}_n \to \mathrm{B\,PU}_n$, where the map $\mathrm{B}\Delta : \mathrm{B}S^1 \to \mathrm{BU}_n$ is that induced by the diagonal. Since $\tilde{\mathrm{H}}^i(\mathrm{B\,PU}_n,\mathbb{Z}) = 0$ for $i \leq 2$, and since $\mathrm{H}^*(\mathrm{B}S^1,\mathbb{Z}) = \mathbb{Z}[t]$ where $|t| = 2$, the Serre exact sequence of low-dimensional terms in the Serre spectral sequence for $\mathrm{B}S^1 \to \mathrm{BU}_n \to \mathrm{B\,PU}_n$ yields an exact sequence

$$0 = \mathrm{H}^3(\mathrm{B}S^1,\mathbb{Z}) \to \mathrm{H}^4(\mathrm{B\,PU}_n,\mathbb{Z}) \xrightarrow{\mathrm{B}\rho^*} \mathrm{H}^4(\mathrm{BU}_n,\mathbb{Z}) \xrightarrow{\mathrm{B}\Delta^*} \mathrm{H}^4(\mathrm{B}S^1,\mathbb{Z})$$

so that the map $\mathrm{B}\rho^*$ is the inclusion $\ker \mathrm{B}\Delta^* \to \mathrm{H}^4(\mathrm{BU}_n,\mathbb{Z})$.

The pull-back along Δ of the universal bundle $\mathrm{EU}_n \to \mathrm{BU}_n$ has total Chern class $(1+t)^n$, from which it follows that $\mathrm{B}\Delta^*(c_1) = nt$, implying that $\mathrm{B}\Delta^*(c_1^2) = n^2t^2$ and $\mathrm{B}\Delta^*(c_2) = \binom{n}{2}t^2$. The description of $\ker \mathrm{B}\Delta^* = \mathrm{H}^4(\mathrm{B\,PU}_n,\mathbb{Z})$ is now elementary. □

Write d for the unique class in $\mathrm{H}^4(\mathrm{B\,PU}_n, \mathbb{Z})$ such that $\mathrm{B}\rho^*(d) = \frac{n-1}{2}c_1^2 - nc_2$ when n is odd, or $\mathrm{B}\rho^*(d) = (n-1)c_1^2 - 2nc_2$ when n is even.

Lemma 4.3. *When $n \geq 2$, there are unique generators $y_1 \in \mathrm{H}^1(\mathrm{U}_n, \mathbb{Z}) \cong \mathbb{Z}$ and $y_3 \in \mathrm{H}^3(\mathrm{U}_n, \mathbb{Z}) \cong \mathbb{Z}$ such that in the cohomology Serre spectral sequence for $\mathrm{U}_n \to \mathrm{EU}_n \to \mathrm{BU}_n$*

$$\begin{aligned} d_2(y_1) &= c_1, & d_3(y_3) &= 0, \\ d_2(y_3) &= 0, & d_4(y_3) &= c_2 \mod c_1^2. \end{aligned}$$

Proof. This follows immediately from the Serre spectral sequence associated with $\mathrm{U}_n \to \mathrm{EU}_n \to \mathrm{BU}_n$. ☐

Lemma 4.4. *The low-degree cohomology of $\mathrm{H}^*(\mathrm{PU}_n, \mathbb{Z})$ is*

$$\begin{aligned} \mathrm{H}^1(\mathrm{PU}_n, \mathbb{Z}) &= 0, & \mathrm{H}^3(\mathrm{PU}_n, \mathbb{Z}) &= \mathbb{Z} \cdot y, \\ \mathrm{H}^2(\mathrm{PU}_n, \mathbb{Z}) &= \mathbb{Z}/n \cdot x, & \mathrm{H}^4(\mathrm{PU}_n, \mathbb{Z}) &= \mathbb{Z}/m \cdot x^2, \end{aligned}$$

where $m = n$ if n is odd and $m = n/2$ if n is even. The class y is uniquely determined by the requirement that $\rho^*(y) = y_3$ when n is odd, $\rho^*(y) = 2y_3$ when n is even.

See [4, Section 4]. The interested reader can work this out directly from the Serre spectral sequence for the fibration $S^1 \xrightarrow{\Delta} \mathrm{U}_n \to \mathrm{PU}_n$.

Proof of Proposition. The integer m is defined as in Lemma 4.4.

The lower-left corner of the Serre spectral sequence associated with $\mathrm{PU}_n \to \mathrm{E\,PU}_n \to \mathrm{B\,PU}_n$ is displayed in figure 1. Since the spectral sequence converges to the cohomology of the contractible space $\mathrm{E\,PU}_n$ we can deduce a great deal about the terms and differentials. In the first place, the d_3-differential $d_3 : (\mathbb{Z}/n) \cdot x \to \mathrm{H}^3(\mathrm{B\,PU}_n, \mathbb{Z})$ is an isomorphism.

$$\begin{tikzcd}
(\mathbb{Z}/m) \cdot x^2 \arrow[ddrrr, "d_3"] & & & & & \\
\mathbb{Z} \cdot y \arrow[drr, "d_2"] \arrow[ddrrrr, bend right, "d_4"] & & & & & \\
(\mathbb{Z}/n) \cdot x \arrow[ddrrr, bend right, "d_3"] & & \mathbb{Z}/n & \mathbb{Z}/n & & \\
0 & & & & & \\
\mathbb{Z} & 0 & 0 & \mathrm{H}^3(\mathrm{B\,PU}_n, \mathbb{Z}) & \mathrm{H}^4(\mathrm{B\,PU}_n, \mathbb{Z}) & \mathrm{H}^5(\mathrm{B\,PU}_n, \mathbb{Z})
\end{tikzcd}$$

FIGURE 1. The E_2-page of the Serre spectral sequence associated with $\mathrm{PU}_n \to \mathrm{E\,PU}_n \to \mathrm{B\,PU}_n$.

In the second place we show that the map labeled d_2 in Figure 1 is surjective. We refer to this map simply as d_2. We deduce the existence of a commutative square

$$\begin{tikzcd}
\ker(d_2) \arrow[r, hook] \arrow[d, two heads, "d_4"] & \mathbb{Z} \cdot y \arrow[r, "\cong"] & \mathbb{Z} \cdot y_3 \arrow[d, "d_4"] \\
\mathrm{H}^4(\mathrm{B\,PU}_n, \mathbb{Z}) \arrow[rr] & & \mathbb{Z} \cdot c_2
\end{tikzcd} \tag{3}$$

where the vertical map on the right is that of the Serre spectral sequence associated to the fiber sequence $\mathrm{U}_n \to \mathrm{EU}_n \to \mathrm{BU}_n$. The subgroup $\ker(d_2)$ is generated by by for some unique integer b such that $d_4(by) = d$ in $\mathrm{H}^4(\mathrm{B\,PU}_n, \mathbb{Z})$, and following the arrows counterclockwise and using Lemma 4.2, we see that by has image $-\epsilon(n)c_2$ in $\mathbb{Z} \cdot c_2$. Following the arrows clockwise, and using Lemmas 4.3 and 4.4, we see that by has image bc_2 in $\mathbb{Z} \cdot c_2$

when n is odd and image $2bc_2$ when n is even. In particular, it follows that $b = -n$. Since $\ker(d_2) = n\mathbb{Z} \cdot y$, we see that d_2 is surjective.

In the third place, $d_3(x^2) = 2xd_3(x)$, so that ax^2 is in the kernel of the d_3-differential if and only if $2a \equiv 0 \pmod{n}$, which is to say that $a \equiv 0 \pmod{m}$ and so the d_3-differential is injective on $\mathbb{Z}/m \cdot x^2$. One may now check directly that there are no nonzero groups which can support a differential $d_i : A \to \mathrm{H}^5(\mathrm{B}\,\mathrm{PU}_n, \mathbb{Z})$, so the group $\mathrm{H}^5(\mathrm{B}\,\mathrm{PU}_n, \mathbb{Z})$ consists of permanent non-zero classes, and since $\mathrm{E}\,\mathrm{PU}_n$ is contractible, it that follows $\mathrm{H}^5(\mathrm{B}\,\mathrm{PU}_n, \mathbb{Z}) = 0$. □

5. THE OBSTRUCTION THEORY OF $\mathrm{B}\,\mathrm{PU}_n$

By definition, the topological index $\mathrm{ind}_{\mathrm{top}}(\alpha)$ is the greatest common divisor of the integers n such that $\alpha \in \mathrm{H}^3(X, \mathbb{Z}) = \mathrm{H}^2(X, S^1)$ is in the image of the connecting homomorphism $\mathrm{H}^1(X, \mathrm{PU}_n) \to \mathrm{H}^2(X, S^1)$. In homotopical terms, it is the GCD of the integers n such that a lift exists in the diagram

$$\begin{tikzcd} & \mathrm{B}\,\mathrm{PU}_n \arrow[d] \\ X \arrow[ur, dashed] \arrow[r, "\alpha"] & K(\mathbb{Z},3). \end{tikzcd} \tag{4}$$

Here the map $\mathrm{B}\,\mathrm{PU}_n \to K(\mathbb{Z}, 3)$ is the application of the classifying-space functor to the map $\mathrm{PU}_n \to \mathrm{B}S^1$ which classifies the quotient map $\mathrm{U}_n \to \mathrm{PU}_n$.

The projective unitary group PU_n is the quotient of U_n by its center, which is the image of a homomorphism $S^1 \to U_n$. The map $S^1 \to \mathrm{U}_n$ of S^1 onto the center of U_n has homotopy type equivalent to n times a generator of $\pi_1\mathrm{U}_n$. Therefore,

$$\pi_i\,\mathrm{PU}_n \cong \begin{cases} \mathbb{Z}/n & \text{if } i = 1, \\ \pi_i\mathrm{U}_n & \text{otherwise.} \end{cases}$$

Bott computed [5, Theorem 5] the following homotopy groups of the unitary groups U_n:

$$\pi_i\mathrm{U}_n \cong \begin{cases} 0 & \text{if } i < 2n \text{ is even,} \\ \mathbb{Z} & \text{if } i < 2n \text{ is odd,} \\ \mathbb{Z}/n! & \text{if } i = 2n. \end{cases}$$

Moreover, the standard inclusion maps $\mathrm{U}_n \to \mathrm{U}_{n+1}$ induce isomorphisms

$$\pi_i\mathrm{U}_n \to \pi_i\mathrm{U}_{n+1}$$

whenever $i < 2n$.

The first two stages of the Postnikov tower of $\mathrm{B}\,\mathrm{PU}_n$ therefore takes the following form

$$\begin{tikzcd} K(\mathbb{Z},4) \arrow[r] & \mathrm{B}\,\mathrm{PU}_n[4] \arrow[d] \\ & \mathrm{B}\,\mathrm{PU}_n[2] \simeq K(\mathbb{Z}/n,2). \end{tikzcd} \tag{5}$$

Lemma 5.1. *The map* $\mathrm{B}\,\mathrm{PU}_n \to K(\mathbb{Z}, 3)$ *of diagram* (4) *can be factored as* $\mathrm{B}\,\mathrm{PU}_n \to \mathrm{B}\,\mathrm{PU}_n[2] \simeq K(\mathbb{Z}/n, 2)$ *and* $\beta : K(\mathbb{Z}/n, 2) \to K(\mathbb{Z}, 3)$.

Proof. This follows from comparing the presentation of PU_n as the quotient of U_n by S^1 and its presentation as the quotient of SU_n by its center, $\mathbb{Z}/n$. □

The map classifying $\mathrm{B}\,\mathrm{PU}_n[4] \to K(\mathbb{Z}/n, 2)$ is a map $f_n : K(\mathbb{Z}/n, 2) \to K(\mathbb{Z}, 5)$, to wit a cohomology class $f_n \in \mathrm{H}^5(K(\mathbb{Z}/n), \mathbb{Z})$. It is to consideration of this class that we now turn.

Lemma 5.2. *The maps* $\pi_i \mathrm{B}\,\mathrm{PU}_n \to \pi_i \mathrm{B}\,\mathrm{PU}_{nr}$ *induced by* r*-fold direct summation are multiplication by* r *in the stable range* $i < 2n+1$.

Proof. Compare with U_n. ☐

When X is a CW complex of cohomological dimension at most 6, then there is a natural bijection

$$[X, \mathrm{B}\,\mathrm{PU}_n] \cong [X, \mathrm{B}\,\mathrm{PU}_n[5]]$$

given by the truncation map $\mathrm{B}\,\mathrm{PU}_n \to \mathrm{B}\,\mathrm{PU}_n[5]$ in the Postnikov tower of $\mathrm{B}\,\mathrm{PU}_n$, where

$$\pi_i \mathrm{B}\,\mathrm{PU}_n[5] = \begin{cases} \pi_i \mathrm{B}\,\mathrm{PU}_n & \text{if } i \leq 5, \\ 0 & \text{otherwise.} \end{cases}$$

Indeed, all obstructions to lifting a map $X \to \mathrm{B}\,\mathrm{PU}_n[5]$ to a map $X \to \mathrm{B}\,\mathrm{PU}_n$ occur in $\mathrm{H}^{k+1}(X, \pi_k \mathrm{B}\,\mathrm{PU}_n)$ where $k \geq 6$. These groups are zero by hypothesis, and the lifts are unique.

The maps $\mathrm{B}\,\mathrm{PU}_n \to \mathrm{B}\,\mathrm{PU}_{nr}$ induced by r-fold direct summation in turn induce maps of the corresponding fibrations above, and in particular a map $K(\mathbb{Z}/n, 2) \to K(\mathbb{Z}/nr, 2)$.

Lemma 5.3. *The pullback map*

$$\mathrm{H}^5(K(\mathbb{Z}/nr, 2), \mathbb{Z}) \to \mathrm{H}^5(K(\mathbb{Z}/n, 2), \mathbb{Z}),$$

takes f_{nr} *to* $r \cdot f_n$ *for all integers* $n > 1$ *and* $r > 0$.

Proof. This follows from the functoriality of Postnikov towers. Indeed, by Lemma 5.2 the map $\pi_i \mathrm{B}\,\mathrm{PU}_n \to \pi_i \mathrm{B}\,\mathrm{PU}_{nr}$ is multiplication by r for all i, as one can see by comparing to $\mathrm{U}_n \to \mathrm{U}_{nr}$. It follows that there is a commutative diagram

$$\begin{tikzcd}
K(\mathbb{Z}/n, 2) \arrow[r, "f_n"] \arrow[d] & K(\mathbb{Z}, 5) \arrow[d] \\
K(\mathbb{Z}/nr, 2) \arrow[r, "f_{nr}"] & K(\mathbb{Z}, 5).
\end{tikzcd}$$

Hence the result. ☐

Recall that we have defined $\epsilon(n)$ to be n if n is odd and $2n$ if n is even.

Proposition 5.4. *The first few integral cohomology groups of* $K(\mathbb{Z}/n, 2)$ *are*

$$\mathrm{H}^1(K(\mathbb{Z}, n), \mathbb{Z}) = \mathrm{H}^2(K(\mathbb{Z}, n), \mathbb{Z}) = \mathrm{H}^4(K(\mathbb{Z}, n), \mathbb{Z}) = 0,$$

and

$$\mathrm{H}^3(K(\mathbb{Z}/n), \mathbb{Z}) = \mathbb{Z}/n \cdot \beta \qquad \mathrm{H}^5(K(\mathbb{Z}/n, 2), \mathbb{Z}) = \mathbb{Z}/\epsilon(n) \cdot Q.$$

Here β *denotes the unreduced Bockstein of* ι*, the fundamental class, and* Q *denotes either* $\beta(\iota^2)$ *if* n *is odd, or a class such that* $2Q = \beta(\iota^2)$ *if* n *is even.*

This may be deduced from the calculations of homology given in [7] or directly from the Serre spectral sequence for $K(\mathbb{Z}/n, 1) \to * \to K(\mathbb{Z}/n, 2)$. In the case where n is even, the class Q is $2n$-torsion, and therefore there exist two classes $P_2 \in \mathrm{H}^4(K(\mathbb{Z}/n, 2), \mathbb{Z}/2n)$ with the property that $\beta(P_2) = Q$. We fix one of these classes and declare it to be P_2. We call this class the *Pontryagin square* of ι, in extension of the ordinary usage of this term in the case when n is a power of 2.

Proposition 5.5. *Let* $n \geq 2$ *be an integer. The element* $f_n \in \mathrm{H}^5(K(\mathbb{Z}/n, 2), \mathbb{Z})$ *is a generator.*

$$\begin{tikzcd}
\mathbb{Z} \arrow[ddddrrrrr, "f_n"] & & & & & \\
0 & & & & & \\
0 & & & & & \\
0 & & & & & \\
\mathbb{Z} & 0 & 0 & \mathbb{Z}/n & 0 & \mathbb{Z}/(\epsilon n)
\end{tikzcd}$$

FIGURE 2. The Serre spectral sequence associated to $\mathrm{B\,PU}_n[4] \to K(\mathbb{Z}/n, 2)$

Proof. Associated to the fibration $\mathrm{B\,PU}_n[4] \to K(\mathbb{Z}/n, 2)$, we have a Serre spectral sequence for integral cohomology, part of the E_2-page of which is illustrated in figure 2. Therefore $\mathrm{H}^5(\mathrm{B\,PU}_n[4], \mathbb{Z}) = \frac{\mathbb{Z}}{\epsilon n} \Big/ \mathrm{im}(f_n)$.

In the case where $n > 2$, because $\pi_5(\mathrm{B\,PU}_n) = 0$, it follows that $\mathrm{B\,PU}_n[4] \simeq \mathrm{B\,PU}_n[5]$, and so $\mathrm{H}^5(\mathrm{B\,PU}_n[4], \mathbb{Z}) = \mathrm{H}^5(\mathrm{B\,PU}_n, \mathbb{Z}) = 0$, and f_n is consequently surjective.

In the case where $p = 2$, we discover another extension problem, associated to the principal fibration $K(\mathbb{Z}/2, 5) \to \mathrm{B\,PU}_2[5] \to \mathrm{B\,PU}_2[4]$. Since $\mathrm{H}^5(\mathrm{B\,PU}_2[5], \mathbb{Z}) = \mathrm{H}^5(\mathrm{B\,PU}_2, \mathbb{Z}) = 0$, there is an exact sequence $0 \to \mathbb{Z}/4 \Big/ \mathrm{im}\, f_2 \to 0 \to \mathbb{Z}/2 \to *$. By examining the part of the Serre spectral sequence appearing in figure 3, it follows that f_2 is surjective as well. □

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[dddddrrrrrr, "d_6"] & & & & & & \\
0 & & & & & & \\
0 & & & & & & \\
0 & & & & & & \\
0 & & & & & & \\
\mathbb{Z} & 0 & 0 & \mathbb{Z}/2 & 0 & \mathbb{Z}/4 \Big/ \mathrm{im}\, f_2 & *
\end{tikzcd}$$

FIGURE 3. The Serre spectral sequence associated to $\mathrm{B\,PU}_2[5] \to \mathrm{B\,PU}_2[4]$

6. RELATING THE OBSTRUCTION THEORY TO THE DESCENT SPECTRAL SEQUENCE

Definition 6.1. Let X be a space, and n a positive integer such that $\mathrm{H}^2(X, \mathbb{Z})$ is n-divisible. Suppose $\alpha \in \mathrm{H}^3(X, \mathbb{Z})$ is an n-torsion class. Then we write $\tilde{\alpha} \in \mathrm{H}^2(X, \mathbb{Z}/n)$ for the unique class such that $\beta(\tilde{\alpha}) = \alpha$.

Proposition 6.2. *Let X be a connected finite CW complex of cohomological dimension $d \leq 5$. Let $\alpha \in \mathrm{H}^3(X, \mathbb{Z})$ be an n-torsion class, and suppose that $\mathrm{H}^2(X, \mathbb{Z}) = 0$. Then $\mathrm{ind}_{\mathrm{top}}(\alpha) = \mathrm{per}_{\mathrm{top}}(\alpha)\, \mathrm{ord}(f_n \circ \tilde{\alpha})$.*

Proof. For any r there exists a unique factorization of α as follows

$$X \longrightarrow K(\mathbb{Z}/nr, 2) \longrightarrow K(\mathbb{Z}, 3).$$

Since X is of cohomological dimension $d \leq 5$, the only obstruction to finding a lift

$$\begin{tikzcd}
 & \mathrm{B\,PU}_{nr} \arrow[d] \\
X \arrow[r] \arrow[ur, dashed] & K(\mathbb{Z}/nr, 2)
\end{tikzcd}$$

is the first one arising in the Postnikov tower of $\mathrm{B\,PU}_{nr}$, as presented in diagram (5) and the subsequent discussion. There is a lift if and only if $X \to K(\mathbb{Z}/nr,2) \xrightarrow{f_{nr}} K(\mathbb{Z},5)$ is nullhomotopic, which holds if and only if $r(f_n \circ \tilde{\alpha}) \simeq 0$ by Lemma 5.3. The result follows. ☐

Corollary 6.3. *Let Y denote $\mathrm{sk}_5(K(\mathbb{Z}/nr,2))$, and let $\beta^Y \in \mathrm{H}^3(Y,\mathbb{Z})$ denote the restriction of the Bockstein β to Y. Then $\mathrm{ind}_{\mathrm{top}}(\beta^Y) = \mathrm{per}_{\mathrm{top}}(\beta^Y)\,\mathrm{ord}(f_n) = \mathrm{per}_{\mathrm{top}}(\beta^Y)\,\mathrm{ord}(G(\beta^Y))$. In particular, $\mathrm{ord}(G(\beta^Y)) = \mathrm{ord}(G(\beta)) = \epsilon(n)$.*

Proof. One combines the proposition with Theorem B and Proposition 5.5. ☐

$$\begin{tikzcd}
\mathbb{Z} & 0 & 0 & \mathbb{Z}/n & 0 & \mathbb{Z}/\epsilon(n) \\
\mathbb{Z} & 0 & 0 & \mathbb{Z}/n & 0 & \mathbb{Z}/\epsilon(n) \\
\mathbb{Z} \arrow[uur, "\smile\beta"] \arrow[uuuuurr, bend right, "d_5^\beta(n)"'] & 0 & 0 & \mathbb{Z}/n & 0 & \mathbb{Z}/\epsilon(n)
\end{tikzcd}$$

FIGURE 4. The descent spectral sequence for $\mathrm{KU}^*(K(\mathbb{Z}/n,2))_\beta$, twisted by the Bockstein.

Theorem 6.4 (Relation (2))**.** *Let X be a space and let n be an integer. Let $\xi \in \mathrm{H}^2(X,\mathbb{Z}/n)$ and let $Q(\xi)$ denote either $\beta_n(\xi^2)$ if n is odd or $Q(\xi) = \beta_{2n}(P_2(\xi))$ if n is even. The class $Q(\xi)$ is $\epsilon(n)$-torsion, and*

$$\left\langle \frac{n}{\mathrm{per}_{\mathrm{top}}(\beta_n(\xi))} G(\beta_n(\xi)) \right\rangle = \langle [Q(\xi)] \rangle$$

in $\mathrm{H}^5(X,\mathbb{Z})/(\mathrm{H}^2(X,\mathbb{Z}) \smile \beta_n(\xi))$.

Proof. We write β for β_n throughout. The statement regarding the order of $Q(\xi)$ is immediate. There is a tautological map $X \xrightarrow{\xi} K(\mathbb{Z}/n,2)$, which induces a map of descent spectral sequences converging conditionally to the map $K^*(K(\mathbb{Z}/n,2))_\beta \to K^*(X)_{\beta(\xi)}$. We consider the differentials supported by $\mathbb{Z} = \mathrm{E}_2^{0,0}$ for both spaces. The d_3-differentials take the form

$$\begin{tikzcd}
\mathbb{Z} \arrow[r, equal] \arrow[d, "d_3^\beta"] & \mathbb{Z} \arrow[d, "d_3^{\beta(\xi)}"] \\
\mathrm{H}^3(K(\mathbb{Z}/n,2),\mathbb{Z}) \arrow[r] & \mathrm{H}^3(X,\mathbb{Z}).
\end{tikzcd}$$

The d_5-differentials take the form

$$\begin{tikzcd}
n\mathbb{Z} \arrow[r, hook] \arrow[d, two heads, "d_5^\beta"] & \mathrm{per}_{\mathrm{top}}(\beta(\xi))\mathbb{Z} \arrow[d, "d_5^{\beta(\xi)}"] \\
\mathbb{Z}/\epsilon(n) \arrow[r] & \mathrm{H}^5(X,\mathbb{Z})/(\mathrm{H}^2(X,\mathbb{Z}) \smile \beta(\xi))
\end{tikzcd}$$

The left vertical map is $n \mapsto G(\beta)$ and is surjective. In particular $d_5^\beta(n) = u'\beta Q$, since Q generates $\mathrm{H}^5(K(\mathbb{Z}/n,2),\mathbb{Z})$. The top horizontal map takes the generator $n \in n\mathbb{Z}$ to $\frac{n}{\mathrm{per}_{\mathrm{top}}(\beta(\xi))}\mathrm{per}_{\mathrm{top}}(\beta(\xi)) \in \mathrm{per}_{\mathrm{top}}(\beta(\xi))\mathbb{Z}$. The bottom horizontal map takes Q tautologically to $Q(\xi)$. The right vertical map therefore takes $\frac{n}{\mathrm{per}_{\mathrm{top}}(\beta(\xi))}\mathrm{per}_{\mathrm{top}}(\beta(\xi))$ to $uQ(\xi)$. The class of $d_5^{\beta(\xi)}(\mathrm{per}_{\mathrm{top}}\,\beta(\xi))$ in $\mathrm{H}^5(X,\mathbb{Z})/(\mathrm{H}^2(X,\mathbb{Z})\smile\beta(\xi))$ is $G(\beta(\xi))$ by definition, so the relation

$$\frac{n}{\mathrm{per}_{\mathrm{top}}(\beta(\xi))}G(\beta(\xi))=Q(\xi)$$

follows. □

If $\alpha\in\mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$ is any class, then there exists some potentially non-unique lift $\xi\in\mathrm{H}^2(X,\mathbb{Z}/n)$ such that $\beta(\xi)=\alpha$. By this method, $G(\alpha)=G(\beta(\xi))$ may be computed, and one deduces further that $G(\alpha)$ is $\epsilon(n)$ torsion, since ξ is n-torsion.

Theorem B. *Let X be a finite CW complex of cohomological dimension at most 6. Then, for any $\alpha\in\mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$,*

$$\mathrm{ind}_{\mathrm{top}}(\alpha)\,\Big|\,\begin{cases}\mathrm{per}_{\mathrm{top}}(\alpha)^2 & \text{if } \mathrm{per}_{\mathrm{top}}(\alpha) \text{ is odd}\\ 2\,\mathrm{per}_{\mathrm{top}}(\alpha)^2 & \text{if } \mathrm{per}_{\mathrm{top}}(\alpha) \text{ is even.}\end{cases}$$

These upper-bounds on the index are the best possible.

Proof. The bounds hold by Theorem A and the observation that $G(\alpha)$ is $\epsilon(n)$-torsion. The bounds are realized by $\beta\in\mathrm{H}^3(\mathrm{sk}_5(K(\mathbb{Z}/n,2)),\mathbb{Z})$, for which $G(\beta)$ has order precisely $\epsilon(n)$ by Corollary 6.3. □

7. SMOOTH COMPLEX 3-FOLDS

In the preceding paragraphs, we have analyzed the topological period-index problem for 6-dimensional CW complexes. Here, we consider the consequences for smooth projective complex 3-folds X. Any such X has the homotopy type of a finite 6-dimensional CW complex. As a result, we find the existence of topological projective bundles on X, and we also outline a potential method for constructing counterexamples to the period-index conjecture for function fields at the prime 2 in dimension 3.

If X is a smooth complex variety and $\alpha\in\mathrm{Br}(X)$, we let $\overline{\alpha}$ denote the image of α under the natural map $\mathrm{Br}(X)\to\mathrm{Br}_{\mathrm{top}}(X)=\mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$.

Theorem 7.1. *Let X be a smooth projective complex 3-fold, and let $\alpha\in\mathrm{Br}(X)$. Then,*

$$\mathrm{ind}_{\mathrm{top}}(\overline{\alpha})\,\Big|\,\begin{cases}\mathrm{per}_{\mathrm{top}}(\overline{\alpha})^2 & \text{if } \mathrm{per}_{\mathrm{top}}(\overline{\alpha}) \text{ is odd}\\ 2\,\mathrm{per}_{\mathrm{top}}(\overline{\alpha})^2 & \text{if } \mathrm{per}_{\mathrm{top}}(\overline{\alpha}) \text{ is even.}\end{cases}$$

If X is moreover simply connected, then $\mathrm{ind}_{\mathrm{top}}(\overline{\alpha})=\mathrm{per}_{\mathrm{top}}(\overline{\alpha})$.

Proof. The first statement is simply an application of Theorem B. The second statement follows from Theorem A and Poincaré duality: if X is simply connected, then $\mathrm{H}^5(X,\mathbb{Z})=0$, so $G(\overline{\alpha})=0$. □

We can use the topological results in another way. Suppose we wanted to disprove the period-index conjecture. A natural approach would be to construct a smooth complex 3-fold X and a class $\overline{\alpha}\in\mathrm{H}^3(X,\mathbb{Z})_{\mathrm{tors}}$ such that $\mathrm{per}_{\mathrm{top}}(\overline{\alpha})=2$ and $\mathrm{ind}_{\mathrm{top}}(\overline{\alpha})=8$. Indeed, given such a class, we could lift it to a class $\alpha\in\mathrm{Br}(X)$ such that $\mathrm{per}(\alpha)=2$. Then,

$$\mathrm{ind}_{\mathrm{top}}(\overline{\alpha})\,|\,\mathrm{ind}(\alpha).$$

Write $\eta\to X$ for the generic point. Since $\mathrm{Br}(X)\to\mathrm{Br}(\eta)$ is injective, $\mathrm{per}(\alpha_\eta)=\mathrm{per}(\alpha)=2$. But, we also have $\mathrm{ind}(\alpha_\eta)=\mathrm{ind}(\alpha)$ by the following proposition, and hence the period-index conjecture would be false.

Proposition 7.2. *Let X be a regular noetherian scheme with function field K, and let $\alpha\in\mathrm{Br}(X)$. Then,* $\mathrm{ind}(\alpha)=\mathrm{ind}(\alpha_K)$.

Proof. The following proof was told to the first author by David Saltman at the AIM workshop "Quadratic forms, division algebras, and patching" in January 2011. Recall that for a given Brauer class α, there is a notion of α-twisted coherent sheaf (see, for instance, [13]). These objects form an abelian category, and we may take the K-theory of this category, obtaining $\mathbf{G}_0^{\alpha}(X)$. Similarly, there is a notion of α-twisted vector bundle, and we can produce K-groups $\mathbf{K}_0^{\alpha}(X)$. The index of α then has the following nice description: it is a generator of the image of the natural rank map $\mathbf{K}_0^{\alpha}(X) \to \mathbb{Z}$. Moreover, when X is regular, we have natural isomorphisms $\mathbf{K}_0^{\alpha}(X) \cong \mathbf{G}_0^{\alpha}(X)$. There are extension by zero maps $j_* : \mathbf{G}_0^{\alpha}(K) \to \mathbf{G}_0^{\alpha}(X)$, where $j : \operatorname{Spec} K \to X$. Under the isomorphisms with K-theory, extension by zero is compatible with the rank map. So, there exists an α-twisted coherent sheaf on X of rank $\operatorname{ind}(\alpha_K)$. Since $\mathbf{G}_0^{\alpha}(X) \cong \mathbf{K}_0^{\alpha}(X)$, we see that

$$\operatorname{ind}(\alpha) \,|\, \operatorname{ind}(\alpha_K).$$

But, we also know in general that

$$\operatorname{ind}(\alpha_K) \,|\, \operatorname{ind}(\alpha),$$

so the result follows. □

Putting everything together, we have proven the following theorem.

Theorem C. *If there exists a smooth projective complex* 3*-fold* X *and a* 2*-torsion class* $\alpha \in \operatorname{Br}(X)$ *such that* $2G(\overline{\alpha}) \neq 0$*, where* $\overline{\alpha}$ *is the image of* α *in* $\mathrm{H}^3(X, \mathbb{Z})_{\mathrm{tors}}$*, then the period-index conjecture is false at the prime* 2 *in dimension* 3*: the image of the class* α *in the function field* $\mathbb{C}(X)$ *has period* 2 *and index at least* 8.

On the other hand, an unpublished result of Daniel Krashen says that if X is a complex 3-fold and $\alpha \in \operatorname{Br}(\mathbb{C}(X))$ has period 2, then $\operatorname{ind}(\alpha)$ divides 8.

If such an example is found, it is natural to ask if an "asymptotic" version of the conjecture is true. That is, the conjecture is true for primes p in dimension d if $p > N_d$ for some integer N_d that depends only on d. In particular, $N_2 = 1$ and N_3 would equal 2. If the period-index fails, does it fail for purely topological reasons?

REFERENCES

[1] B. Antieau and B. Williams, *The period-index problem for twisted topological K-theory*, ArXiv e-prints (2011), available at `1104.4654`.

[2] M. Atiyah and G. Segal, *Twisted K-theory*, Ukr. Mat. Visn. **1** (2004), no. 3, 287–330; English transl., Ukr. Math. Bull. **1** (2004), no. 3, 291–334.

[3] ———, *Twisted K-theory and cohomology*, Inspired by S. S. Chern, Nankai Tracts Math., vol. 11, World Sci. Publ., Hackensack, NJ, 2006, pp. 5–43.

[4] P. F. Baum and W. Browder, *The cohomology of quotients of classical groups*, Topology **3** (1965), 305–336.

[5] R. Bott, *The space of loops on a Lie group*, Michigan Math. J. **5** (1958), 35–61.

[6] E. H. Brown Jr., *The cohomology of* BSO_n *and* BO_n *with integer coefficients*, Proc. Amer. Math. Soc. **85** (1982), no. 2, 283–288.

[7] H. Cartan, *Détermination des algèbres* $\mathrm{H}_*(\pi, n; \mathbb{Z})$, Séminaire H. Cartan **7** (1954/1955), no. 1, 11-01–11-24.

[8] J.-L. Colliot-Thélène, *Exposant et indice d'algèbres simples centrales non ramifiées*, Enseign. Math. (2) **48** (2002), no. 1-2, 127–146. With an appendix by Ofer Gabber.

[9] A. J. de Jong, *The period-index problem for the Brauer group of an algebraic surface*, Duke Math. J. **123** (2004), no. 1, 71–94.

[10] J. Starr and J. de Jong, *Almost proper GIT-stacks and discriminant avoidance*, Doc. Math. **15** (2010), 957–972.

[11] A. Grothendieck, *Le groupe de Brauer. I. Algèbres d'Azumaya et interprétations diverses*, Séminaire Bourbaki, Vol. 9, Soc. Math. France, Paris, 1995, pp. Exp. No. 290, 199–219.

[12] M. Kameko and N. Yagita, *The Brown-Peterson cohomology of the classifying spaces of the projective unitary groups* $\mathrm{PU}(p)$ *and exceptional Lie groups*, Trans. Amer. Math. Soc. **360** (2008), no. 5, 2265–2284.

[13] M. Lieblich, *Twisted sheaves and the period-index problem*, Compos. Math. **144** (2008), no. 1, 1–31.

[14] J. McCleary, *A user's guide to spectral sequences*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 58, Cambridge University Press, Cambridge, 2001.

[15] J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J., 1974. Annals of Mathematics Studies, No. 76.

[16] A. Vistoli, *On the cohomology and the Chow ring of the classifying space of* PGL_p, J. Reine Angew. Math. **610** (2007), 181–227.

[17] L. M. Woodward, *The classification of principal* PU_n*-bundles over a* 4*-complex*, J. London Math. Soc. (2) **25** (1982), no. 3, 513–524.

UCLA, DEPARTMENT OF MATHEMATICS, 520 PORTOLA PLAZA, LOS ANGELES, CA 90095-1555
E-mail address: `antieau@math.ucla.edu`

USC, DEPARTMENT OF MATHEMATICS, 3620 SOUTH VERMONT AVENUE, LOS ANGELES, CA 90089-2532
E-mail address: `tbwillia@usc.edu`